

hep-ph/0204030
KUNS-1777
BA-02-15
February 7, 2008

Supersymmetric $SU(3)^3$ unification with Anomalous $U(1)_A$ Gauge Symmetry

Nobuhiro Maekawa^a and Qaisar Shafi^b

^a*Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

^b*Bartol Research Institute, University of Delaware, Newark, DE19716, USA*

Abstract

We consider supersymmetric unification based on the gauge symmetry $SU(3)_C \times SU(3)_L \times SU(3)_R$ supplemented by an anomalous $U(1)_A$ gauge symmetry. Realistic fermion masses and mixings are realized, including large mixings in the neutrino sector. We also consider the supersymmetric flavor problem, gauge coupling unification and proton decay. The dominant proton decay mode is expected to be $p \rightarrow e^+ \pi^0$ and the lifetime is estimated to be $\sim 10^{34} - 10^{35}$ years.

^ae-mail: maekawa@gauge.scphys.kyoto-u.ac.jp

^b email: shafi@bartol.udel.edu

1 Introduction

Supersymmetric grand unified theories offer a particularly elegant scenario for unifying the strong and electroweak interactions. Recently, realistic models based on $SO(10)$ [1, 2, 3] and E_6 [4, 5] have been proposed in which an anomalous $U(1)_A$ gauge symmetry[6] plays a critical role. The anomaly is cancelled via the Green Schwarz mechanism[7], and the resulting phenomenology has several attractive features. In particular, all interactions allowed by the symmetries are included in the discussion (there can be undetermined order unity coefficients accompanying the interactions). These models naturally resolve the doublet- triplet problem [8, 9] using the mechanism of reference[10] (see also[11, 12, 13]). Realistic pattern of quark and lepton mass matrices, including large neutrino mixings[14], are realized, using the Froggatt- Nielsen (FN) mechanism[15]. The anomalous $U(1)_A$ also helps explain the hierarchical symmetry breaking scales and the masses acquired by the superheavy particles, and leads precisely to the minimal supersymmetric standard model (MSSM) at low energies. Even though the gauge couplings unify in these schemes slightly below the usual GUT scale 2×10^{16} GeV, dimension five proton decay is sufficiently suppressed, and the decay $p \rightarrow e + \pi$ via gauge mediated dimension six operators may be seen in the near future. Finally, in these models the cutoff scale is lower than the Planck scale M_{Planck} and the μ problem is also resolved.

However, these models require two adjoint Higgs fields to realize DT splitting, which is not so easily realized in the framework of superstring models. In this paper we examine the application of the above approach to grand unified theories with semi-simple unification whose symmetry breaking to MSSM (the minimal supersymmetric standard model) does not require adjoint Higgs field. A particularly attractive example is provided by the gauge symmetry $SU(3)^3 \equiv SU(3)_C \times SU(3)_L \times SU(3)_R$ which is a maximal subgroup of E_6 , and which arises as an effective four dimensional symmetry from the compactification of the $E_8 \times E_8$ heterotic superstring theory on a Calabi-Yau manifold[16]. Phenomenology based on $SU(3)^3$ has been extensively discussed in the past[17, 18, 19]. Our goal here is to apply the techniques of [1]-[5] to gauge symmetry $SU(3)^3$ and elucidate the most important consequences. In particular, we will show how realistic fermion masses and mixings are obtained, including large mixings in the neutrino sector (with the exception of U_{e3}). We also consider SUSY breaking and flavor changing neutral currents, gauge coupling unification and proton decay. While dimension five proton decay turns out to be heavily suppressed, dimension six operators yield a proton lifetime of $\sim 10^{34} - 10^{35}$ years for the decay channel $p \rightarrow e^+ \pi^0$.

2 Matter sector

The matter sector has essentially the same structure as the E_6 model [4], with the **27** of E_6 given in terms of $SU(3)_C \times SU(3)_L \times SU(3)_R$ as

$$\mathbf{27} \rightarrow (\mathbf{3}, \bar{\mathbf{3}}, \mathbf{1}) + (\bar{\mathbf{3}}, \mathbf{1}, \mathbf{3}) + (\mathbf{1}, \mathbf{3}, \bar{\mathbf{3}}). \quad (2.1)$$

Three **27**-plets Ψ_i ($i = 1, 2, 3$) are introduced, and the Yukawa interactions contain appropriate powers of the VEV of FN field $\langle \Theta \rangle = \lambda \Lambda$, which has an anomalous $U(1)_A$ charge $\theta = -1$, namely

$$\lambda^{\psi_i + \psi_j + \phi} \Psi_i \Psi_j \Phi. \quad (2.2)$$

Here Φ is a Higgs field, and throughout this paper we use units in which the cutoff $\Lambda = 1$, and denote all superfields by uppercase letters and their anomalous $U(1)_A$ charges by the corresponding lowercase letters. Using the definitions of the fields $Q(\mathbf{3}, \mathbf{2})_{\frac{1}{6}}$, $U^c(\bar{\mathbf{3}}, \mathbf{1})_{-\frac{2}{3}}$, $D^c(\bar{\mathbf{3}}, \mathbf{1})_{\frac{1}{3}}$, $L(\mathbf{1}, \mathbf{2})_{-\frac{1}{2}}$, $E^c(\mathbf{1}, \mathbf{1})_1$, $N^c(\mathbf{1}, \mathbf{1})_0$, $L'(\mathbf{1}, \mathbf{2})_{\frac{1}{2}}$, $L'(\mathbf{1}, \mathbf{2})_{-\frac{1}{2}}$, $D'(\mathbf{3}, \mathbf{1})_{-\frac{1}{3}}$, $D'(\bar{\mathbf{3}}, \mathbf{1})_{\frac{1}{3}}$, $S(\mathbf{1}, \mathbf{1})$, and their conjugate fields under the standard model (SM) gauge symmetry, the fields $(\mathbf{3}, \bar{\mathbf{3}}, \mathbf{1})$, $(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{3})$, and $(\mathbf{1}, \mathbf{3}, \bar{\mathbf{3}})$ under $SU(3)_C \times SU(3)_L \times SU(3)_R$ are

$$(\mathbf{3}, \bar{\mathbf{3}}, \mathbf{1}) \rightarrow Q + D' \quad (2.3)$$

$$(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{3}) \rightarrow U^c + D^c + D' \quad (2.4)$$

$$(\mathbf{1}, \mathbf{3}, \bar{\mathbf{3}}) \rightarrow L + E^c + N^c + L' + L' + S. \quad (2.5)$$

For future reference, under the breaking $E_6 \rightarrow SO(10)$,

$$\mathbf{27} \rightarrow \underbrace{[Q + U^c + E^c + D^c + L + N^c]}_{\mathbf{16}} + \underbrace{[L' + L' + D' + D']}_{\mathbf{10}} + \underbrace{S}_{\mathbf{1}}. \quad (2.6)$$

Since $D'(L')$ can acquire superheavy mass by combining with a linear combination of D^c and D' (L and L') after the breaking $SU(3)_C \times SU(3)_L \times SU(3)_R$ to the SM gauge group, the remaining massless fields form the three generation matter content of MSSM. Since the Yukawa couplings are determined mainly by the anomalous $U(1)_A$ charges of the massless fields, we would like to know which of the fields among D_i^c and D_i' (L_i and L'_i) ($i = 1, 2, 3$) are massless.

To this end, we discuss how $SU(3)_C \times SU(3)_L \times SU(3)_R$ breaks to MSSM. We introduce the Higgs fields with non-vanishing VEVs $\Phi(\mathbf{1}, \mathbf{3}, \bar{\mathbf{3}})$, $\bar{\Phi}(\mathbf{1}, \bar{\mathbf{3}}, \mathbf{3})$, $C(\mathbf{1}, \mathbf{3}, \bar{\mathbf{3}})$, and $\bar{C}(\mathbf{1}, \bar{\mathbf{3}}, \mathbf{3})$. The VEVs $|\langle \Phi \rangle| = |\langle \bar{\Phi} \rangle| \sim \lambda^{-(\phi + \bar{\phi})/2}$ break $SU(3)_C \times SU(3)_L \times SU(3)_R$ to $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$, while $|\langle C \rangle| = |\langle \bar{C} \rangle| \sim \lambda^{-(c + \bar{c})/2}$ break $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ to the SM gauge group. Here the VEVs are determined by the anomalous $U(1)_A$ charges, and the reason is roughly as follows (We will explain how to determine the VEVs later):

1. Since the interactions are determined by the anomalous $U(1)_A$ charges, the VEV of the gauge invariant operator O with negative charge o is determined as $\langle O \rangle \sim \lambda^{-o}$.
2. For $O = \bar{\Phi}\Phi$, the VEV becomes $\langle \bar{\Phi}\Phi \rangle \sim \lambda^{-(\phi+\bar{\phi})}$.
3. Since the D -flatness condition requires $|\langle \bar{\Phi} \rangle| = |\langle \Phi \rangle|$, we obtain $|\langle \bar{\Phi} \rangle| = |\langle \Phi \rangle| \sim \lambda^{-\frac{1}{2}(\bar{\phi}+\phi)}$.

The massless modes can be determined from the superpotential

$$W_1 = \lambda^{\psi_i+\psi_j+\phi}\Psi_i\Psi_j\Phi + \lambda^{\psi_i+\psi_j+c}\Psi_i\Psi_jC, \quad (2.7)$$

and the VEVs $\langle \Phi \rangle \sim \lambda^{-\frac{1}{2}(\phi+\bar{\phi})}$ and $\langle C \rangle \sim \lambda^{-\frac{1}{2}(c+\bar{c})}$. Here, for simplicity, we have assumed the E_6 like charge assignment in the matter sector, but in principle, we can assign these charges without respecting E_6 symmetry. The mass matrices of $D^{c(l)}$ and D' ($L^{(l)}$ and L') are obtained from

$$M_I = \begin{pmatrix} I_1 & I_2 & I_3 & I'_1 & I'_2 & I'_3 \\ I_1^{c'} & I_2^{c'} & I_3^{c'} & I_1^{c'} & I_2^{c'} & I_3^{c'} \\ \bar{I}_2' & \bar{I}_3' & \bar{I}_1' & \bar{I}_2' & \bar{I}_3' & \bar{I}_1' \end{pmatrix} \lambda^{\frac{1}{2}(\phi-\bar{\phi})}, \quad (2.8)$$

where $I = L, D^c$ and we have defined the parameter r as

$$r \equiv \frac{1}{2}(c - \bar{c} - \phi + \bar{\phi}), \quad (2.9)$$

which we use frequently in the following discussion. Note that the mass matrices are determined by the anomalous $U(1)_A$ charges. Therefore the massless modes are also determined by the charges. As discussed in Ref.[4], as long as we neglect the cases with vanishing coefficients arising from some SUSY based mechanism, the main components of the massless modes can be obtained as follows:

1. $\psi_1 - \psi_3 < r : (I_1, I_2, I_3)$.
2. $0 < r < \psi_1 - \psi_3 : (I_1, I'_1, I_2)$.
3. $\psi_3 - \psi_1 < r < 0 : (I_1, I'_1, I'_2)$.
4. $r < \psi_3 - \psi_1 : (I'_1, I'_2, I'_3)$.

The case (I_1, I'_1, I_2) is interesting, because bi-large neutrino mixing angles can be realized without $\tan\beta$ too small, if we take account of the mixing of subcomponents. Indeed, the massless modes (I_1^0, I_2^0, I_3^0) are given by

$$I_1^0 = I_1 + \lambda^{\psi_1-\psi_3}I_3 + \lambda^{\psi_1-\psi_2+r}I'_2 + \lambda^{\psi_1-\psi_3+r}I'_3, \quad (2.10)$$

$$I_2^0 = I'_1 + \lambda^{\psi_1-\psi_3-r}I_3 + \lambda^{\psi_1-\psi_2}I'_2 + \lambda^{\psi_1-\psi_3}I'_3, \quad (2.11)$$

$$I_3^0 = I_2 + \lambda^{\psi_2-\psi_3}I_3 + \lambda^rI'_2 + \lambda^{\psi_2-\psi_3+r}I'_3, \quad (2.12)$$

where the first terms on the right-hand sides are the main components of these massless modes, and the other terms represent mixing with the other states, I_3 , I'_2 and I'_3 .

The mass matrices for quarks and leptons are obtained from the superpotential

$$W_2 = \lambda^{\psi_i + \psi_j + h} \Psi_i \Psi_j H, \quad (2.13)$$

where $H(\mathbf{1}, \mathbf{3}, \bar{\mathbf{3}})$ contains the MSSM Higgs doublets. If we adopt the charges $\psi_1 = 3 + n$, $\psi_2 = 2 + n$, $\psi_3 = n$, and $h = -2n$, the mass matrices are given by

$$M_U = \begin{pmatrix} U_1^c & U_2^c & U_3^c \\ Q_1 & Q_2 & Q_3 \end{pmatrix} \begin{pmatrix} \lambda^6 & \lambda^5 & \lambda^3 \\ \lambda^5 & \lambda^4 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{pmatrix} \langle L^c(H) \rangle, \quad (2.14)$$

$$M_D(M_E^T \eta^{-1}) = \begin{pmatrix} D_1^{c0}(L_1^0) & D_2^{c0}(L_2^0) & D_3^{c0}(L_3^0) \\ Q_1(E_1^c) & Q_2(E_2^c) & Q_3(E_3^c) \end{pmatrix} \begin{pmatrix} \lambda^6 & \lambda^{6-r} & \lambda^5 \\ \lambda^5 & \lambda^{5-r} & \lambda^4 \\ \lambda^3 & \lambda^{3-r} & \lambda^2 \end{pmatrix} \langle L'(H) \rangle. \quad (2.15)$$

Here $\eta \sim 2-3$ is the renormalization group factor. Then we can obtain the CKM matrix as

$$U_{\text{CKM}} = \begin{pmatrix} 1 & \lambda & \lambda^3 \\ \lambda & 1 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{pmatrix}, \quad (2.16)$$

which reproduces the experimental results if we take $\lambda \sim 0.2$. Since the ratio of the Yukawa couplings of the top and bottom quarks is λ^2 , a small value of $\tan \beta \sim m_t/m_b \cdot \lambda^2$ is predicted by these mass matrices.

The Dirac neutrino mass matrix is given by the 3×6 matrix

$$\begin{pmatrix} S_1 & S_2 & S_3 & N_1^c & N_2^c & N_3^c \\ L_1^0 & L_2^0 & L_3^0 \end{pmatrix} \begin{pmatrix} \lambda^{r+6} & \lambda^{r+5} & \lambda^{r+3} & \lambda^6 & \lambda^5 & \lambda^3 \\ \lambda^6 & \lambda^5 & \lambda^3 & \lambda^{6-r} & \lambda^{5-r} & \lambda^{3-r} \\ \lambda^{r+5} & \lambda^{r+4} & \lambda^{r+2} & \lambda^5 & \lambda^4 & \lambda^2 \end{pmatrix} \langle L^c(H) \rangle \eta, \quad (2.17)$$

which we simply express as

$$M_N = \begin{pmatrix} \lambda^{r+2} & \lambda^2 \end{pmatrix} \otimes \begin{pmatrix} \lambda^4 & \lambda^3 & \lambda \\ \lambda^{4-r} & \lambda^{3-r} & \lambda^{1-r} \\ \lambda^3 & \lambda^2 & 1 \end{pmatrix} \langle L^c(H) \rangle \eta. \quad (2.18)$$

The 6×6 matrix for the right-handed neutrinos ($S_i, i = 1, 2, 3$, and $N_k^c, k = 1, 2, 3$) is obtained as

$$M_R = \lambda^{\psi_i + \psi_j + 2\bar{\phi}} S_i S_j \langle \bar{\Phi} \rangle^2 + \lambda^{\psi_i + \psi_k + \bar{c} + \bar{\phi}} S_i N_k^c \langle \bar{\Phi} \rangle \langle \bar{C} \rangle,$$

$$+\lambda^{\psi_k+\psi_m+2\bar{c}}N_k^cN_m^c\langle\bar{C}\rangle^2 \quad (2.19)$$

$$= \lambda^{2n} \begin{pmatrix} \lambda^{\bar{\phi}-\phi} & \lambda^{(\bar{\phi}-\phi+\bar{c}-c)/2} \\ \lambda^{(\bar{\phi}-\phi+\bar{c}-c)/2} & \lambda^{\bar{c}-c} \end{pmatrix} \otimes \begin{pmatrix} \lambda^6 & \lambda^5 & \lambda^3 \\ \lambda^5 & \lambda^4 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{pmatrix} \quad (2.20)$$

from the interactions

$$\lambda^{\psi_i+\psi_j+2\bar{\phi}}\Psi_i\Psi_j\bar{\Phi}\bar{\Phi} + \lambda^{\psi_i+\psi_j+\bar{c}+\bar{\phi}}\Psi_i\Psi_j\bar{\Phi}\bar{C} + \lambda^{\psi_i+\psi_j+2\bar{c}}\Psi_i\Psi_j\bar{C}\bar{C}, \quad (2.21)$$

by developing the VEVs $\langle\bar{\Phi}\rangle \sim \lambda^{-\frac{1}{2}(\phi+\bar{\phi})}$ and $\langle\bar{C}\rangle \sim \lambda^{-\frac{1}{2}(c+\bar{c})}$. Using the seesaw mechanism [20], we obtain the neutrino mass matrix

$$M_\nu = M_N M_R^{-1} M_N^T = \lambda^{4-2n+c-\bar{c}} \begin{pmatrix} \lambda^2 & \lambda^{2-r} & \lambda \\ \lambda^{2-r} & \lambda^{2-2r} & \lambda^{1-r} \\ \lambda & \lambda^{1-r} & 1 \end{pmatrix} \langle L^c(H) \rangle^2 \eta^2. \quad (2.22)$$

Then, we finally obtain the Maki-Nakagawa-Sakata matrix as

$$U_{\text{MNS}} = \begin{pmatrix} 1 & \lambda^r & \lambda \\ \lambda^r & 1 & \lambda^{1-r} \\ \lambda & \lambda^{1-r} & 1 \end{pmatrix}. \quad (2.23)$$

If we adopt $r = 1/2$, namely,

$$c - \bar{c} = \phi - \bar{\phi} + 1, \quad (2.24)$$

bi-large neutrino mixing angles are realized because of $\lambda^{1/2} \sim 0.5$ as within the $SO(10)$ model in Ref. [1] and E_6 model in Ref. [4]. Moreover, it predicts $U_{e3} \sim \lambda$. Future experiments can see whether there is such a large U_{e3} just below the CHOOZ upper limit $U_{e3} \leq 0.15$ [21] or not. For the neutrino masses, the model predicts $m_{\nu_\mu}/m_{\nu_\tau} \sim \lambda$, which is consistent with the most probable LMA MSW solution for the solar neutrino puzzle. [14]

If we define

$$l \equiv \bar{\phi} - \phi + 2n - 10, \quad (2.25)$$

the neutrino mass matrix is given by

$$M_\nu = \lambda^{-(5+l)} \begin{pmatrix} \lambda^2 & \lambda^{1.5} & \lambda \\ \lambda^{1.5} & \lambda & \lambda^{0.5} \\ \lambda & \lambda^{0.5} & 1 \end{pmatrix} \langle L^c(H) \rangle^2 \eta^2, \quad (2.26)$$

where we have used the relation (2.24). The paramter l can be determined by

$$\lambda^l = \lambda^{-5} \frac{\langle L^c(H) \rangle^2 \eta^2}{m_{\nu_\tau} \Lambda}. \quad (2.27)$$

We are supposing that the cutoff scale Λ is in the range $5 \times 10^{15} \text{ GeV} < \Lambda < 10^{20} \text{ GeV}$, which allows $-3 \leq l \leq 2$. If we choose $l = -2$, the neutrino masses are given by $m_{\nu_\tau} \sim \lambda^{-3} \langle L^c(H) \rangle^2 \eta^2 / \Lambda \sim m_{\nu_\mu} / \lambda \sim m_{\nu_e} / \lambda^2$. If we take $\eta \langle L^c(H) \rangle = 100 \text{ GeV}$, $\Lambda \sim 2 \times 10^{16} \text{ GeV}$ and $\lambda = 0.2$, then we get $m_{\nu_\tau} \sim 6 \times 10^{-2} \text{ eV}$, $m_{\nu_\mu} \sim 1 \times 10^{-2} \text{ eV}$ and $m_{\nu_e} \sim 2 \times 10^{-3} \text{ eV}$. These values are pretty much consistent with the experimental data for atmospheric neutrinos and with the large mixing angle (LMA) MSW solution for the solar neutrino problem.[22]

3 SUSY breaking and FCNC

Let us now discuss SUSY breaking. Since the anomalous $U(1)_A$ charges depend on flavor to produce the hierarchy of Yukawa couplings, generically non-degenerate scalar fermion masses are induced through the anomalous $U(1)_A$ D -term. Under the E_6 -like charge assignment of the matter sector, the SUSY contribution to $K^0 - \bar{K}^0$ mixing is naturally suppressed as in E_6 GUT scenario. The essential point is that the anomalous $U(1)_A$ charge of D_1^{c0} becomes the same as that of D_2^{c0} , because the field $D_1^{c0} \sim D_1^c$ and $D_2^{c0} \sim D_1^c$ arise from a single field Ψ_1 of E_6 . Since the constraints from $K^0 - \bar{K}^0$ mixing to the ratio $\delta \equiv \Delta / \tilde{m}^2$, where Δ is the mixing of sfermion mass matrices and $\tilde{m}$ is the average of sfermion mass, are

$$\sqrt{|\text{Re}(\delta_{12}^D)_{LL}(\delta_{12}^D)_{RR}|} \leq 2.8 \times 10^{-3} \left(\frac{\tilde{m}_q(\text{GeV})}{500} \right), \quad (3.1)$$

$$|\text{Re}(\delta_{12}^D)_{LL}|, |\text{Re}(\delta_{12}^D)_{RR}| \leq 4.0 \times 10^{-2} \left(\frac{\tilde{m}_q(\text{GeV})}{500} \right), \quad (3.2)$$

and the former constraint is much stronger than the latter constraint, suppression of $(\delta_{12}^D)_{RR}$ makes the constraint on the SUSY breaking sector much weaker. Here we use the notation $(\delta_{ij}^F)_{XY}$, where $F = U, D, N, E$, the chirality index is $X, Y = L, R$, and the generation index is $i, j = 1, 2, 3$, as defined in Ref. [25]. As in the usual anomalous $U(1)_A$ scenario, Δ can be estimated as

$$(\Delta_{ij}^F)_{XX} \sim \lambda^{|f_i - f_j|} (|f_i - f_j|) \langle D_A \rangle. \quad (3.3)$$

Therefore, the coincidence of the anomalous $U(1)_A$ charges of D_1^{c0} and D_2^{c0} leads to the suppression of $(\Delta_{12}^D)_{RR}$. This weakens the constraints from $K^0 - \bar{K}^0$ mixing.

To see how weak the constraints become in our scenario, we fix the SUSY breaking sector as follows. At the cutoff scale, we adopt the common gaugino mass $M_{1/2}$ and D -term of anomalous $U(1)_A$ gauge symmetry D_A as non-vanishing SUSY breaking parameters. Then, the scalar fermion mass squared at low energy scales is estimated as

$$\tilde{m}_{F_i}^2 \sim f_i R M_{1/2}^2 + \eta_F M_{1/2}^2, \quad (3.4)$$

where η_F is a renormalization group factor and

$$R \equiv \frac{\langle D_A \rangle}{M_{1/2}^2}. \quad (3.5)$$

The constraint (3.2) for $(\delta_{12}^D)_{LL}$ is rewritten

$$M_{1/2} \geq 1.25 \times 10^4 \lambda \frac{(\psi_1 - \psi_2)R}{(\eta_{D_L} + \frac{\psi_1 + \psi_2}{2}R)^{3/2}} (\text{GeV}). \quad (3.6)$$

Though the main contribution to $(\delta_{12}^D)_{RR}$ vanishes, through the mixing in Eqs. (2.10) and (2.11), $(\delta_{12}^D)_{RR}$ is estimated as

$$(\delta_{12}^D)_{RR} \sim \lambda^{\frac{1}{2}} \frac{\lambda^2(\psi_1 - \psi_2)R}{\eta_{D_R} + \psi_1 R}. \quad (3.7)$$

From Eq. (3.1) for $\sqrt{(\delta_{12}^D)_{LL}(\delta_{12}^D)_{RR}}$, the constraint on the gaugino mass $M_{1/2}$ is given by

$$M_{1/2} \geq 1.8 \times 10^5 \frac{\lambda^{1.75} R(\psi_1 - \psi_2)}{(\eta_D + \psi_1 R)^{1.5}}. \quad (3.8)$$

On the other hand, the $\mu \rightarrow e\gamma$ process gives

$$|(\delta_{12}^E)_{LL}|, |(\delta_{12}^E)_{RR}| \leq 3.8 \times 10^{-3} \left(\frac{\tilde{m}_l (\text{GeV})}{100} \right)^2, \quad (3.9)$$

where $\tilde{m}_l$ is the average mass of the scalar leptons. This constraint is rewritten

$$M_{1/2} \geq 1.6 \times 10^3 \frac{(\lambda(\psi_1 - \psi_2)R)^{1/2}}{\eta_{E_R} + \frac{\psi_1 + \psi_2}{2}R} (\text{GeV}). \quad (3.10)$$

Taking the values $\psi_1 = 9/2$, $\psi_2 = 7/2$, $\eta_{D_L} \sim \eta_{D_R} \sim 6$ and $\eta_{E_R} \sim 0.15$, the rough lower limits on the gaugino mass are in Table I.

Table I. Lower bound on gaugino mass $M_{1/2}$ at the GUT scale (in GeV).

R	0.1	0.3	0.5	1	2
$(\delta_{12}^D)_{LL}$	17	43	61	87	105
$\sqrt{(\delta_{12}^D)_{LL}(\delta_{12}^D)_{RR}}$	78	191	268	373	437
$ (\delta_{12}^E)_{RR} $	431	304	221	161	116

Note that in some range of R , the $\mu \rightarrow e\gamma$ process gives the severest constraint among the FCNC processes.[26] Therefore, the lepton flavour violating processes[26, 27] might be seen in the near future.

4 Higgs sector

In addition to the Higgs with non-vanishing VEVs Φ , $\bar{\Phi}$, C , and $\bar{C}$, we introduce $C''(\mathbf{1}, \mathbf{3}, \bar{\mathbf{3}})$, $\bar{C}''(\mathbf{1}, \bar{\mathbf{3}}, \mathbf{3})$ with vanishing VEVs and several singlets S and Z in order to give superheavy masses to these Higgs fields. The Higgs content is

Table II. The typical values of anomalous $U(1)_A$ charges are listed.

	non-vanishing VEV	vanishing VEV
$(\mathbf{1}, \mathbf{3}, \mathbf{\bar{3}})$	$\Phi(\phi = -3) \ C(c = -3)$	$C'(c' = 0)$
$(\mathbf{1}, \mathbf{\bar{3}}, \mathbf{3})$	$\bar{\Phi}(\bar{\phi} = 2) \ \bar{C}(\bar{c} = 1)$	$\bar{C}'(\bar{c}' = 4)$
$\mathbf{1}$	$Z_i(z_i = -1)(i = 1, 2, 3)$	
$(\mathbf{3}, \mathbf{\bar{3}}, \mathbf{1})$		$Q_L(q_l = 1)$
$(\mathbf{\bar{3}}, \mathbf{3}, \mathbf{1})$		$\bar{Q}_L(\bar{q}_l = 0)$
$(\mathbf{\bar{3}}, \mathbf{1}, \mathbf{3})$		$Q_R(q_r = 1)$
$(\mathbf{3}, \mathbf{1}, \mathbf{\bar{3}})$		$\bar{Q}_R(\bar{q}_r = 0)$

Here the Higgs field H is contained in Φ as in the E_6 case. The Higgs fields Q_L , $\bar{Q}_L$, Q_R , and $\bar{Q}_R$ are introduced only for realizing the same Kac-Moody levels of the three $SU(3)$ gauge groups and they do not play any other role in the following argument.

In this model, the singlet composite operator $\bar{\Phi}\Phi$ plays the same role as the FN field Θ . The D -flatness condition for the anomalous $U(1)_A$ gauge symmetry is

$$D_A = g_A (\xi^2 + \phi|\Phi|^2 + \bar{\phi}|\bar{\Phi}|^2) = 0, \quad (4.1)$$

where ξ^2 is the parameter of the Fayet-Illiopoulos D -term. Since the D -flatness conditions of $SU(3)_L$ and $SU(3)_R$ require $|\langle\Phi\rangle| = |\langle\bar{\Phi}\rangle|$, the D -flatness condition for the anomalous $U(1)_A$ gauge symmetry is rewritten

$$D_A = g_A (\xi^2 + (\phi + \bar{\phi})|\Phi|^2) = 0. \quad (4.2)$$

Thus we obtain $\xi^2 + (\phi + \bar{\phi})|\Phi|^2 = 0$, namely, $|\langle\Phi\rangle| = |\langle\bar{\Phi}\rangle| = \xi$. In this case, since $\bar{\Phi}\Phi$ plays the same role as Θ , the unit of hierarchy becomes $\langle\bar{\Phi}\Phi\rangle = \lambda \sim \xi^2$, which is different from the usual case in which the FN field is just a singlet field Θ and $\langle\Theta\rangle = \xi$. It means that even if ξ has a milder hierarchy, the unit of hierarchy becomes stronger. Using gauge rotation and D -flatness condition for $SU(3)_L \times SU(3)_R$ gauge symmetry, the VEV can be taken as

$$|\langle\Phi\rangle| = |\langle\bar{\Phi}\rangle| = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \lambda^{-\frac{1}{2}(\phi+\bar{\phi})} \end{pmatrix}, \quad (4.3)$$

which breaks $SU(3)_C \times SU(3)_L \times SU(3)_R$ into $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$. In order to determine the VEVs of the other Higgs fields, we examine the following superpotential

$$W = W_{C'} + W_{\bar{C}'} + W_{NV}, \quad (4.4)$$

where W_X denotes the terms linear in the field X , which has vanishing VEV, and W_{NV} includes only the fields with non-vanishing VEVs. From the superpotential

$$W_{NV} = \bar{\Phi}^3 + \bar{\Phi}^2 \bar{C}, \quad (4.5)$$

The fields E_C^c and $\bar{E}_C^c$ are eaten by the Higgs mechanism in breaking $SU(2)_R \times U(1)_{B-L}$ into $U(1)_Y$. The mass spectrum of the remaining fields is $\lambda^{c'+\phi}$ and $\lambda^{c'+\bar{\phi}}$.

The mass matrix M_{D^c} for the fields $D^c, D^{c'}, \bar{D}^c, \bar{D}^{c'}$ is obtained from the interactions

$$\begin{matrix} D_{Q_R}^c \\ D_{Q_R}^{c'} \\ \bar{D}_{Q_L}^{c'} \end{matrix} \begin{pmatrix} \bar{D}_{Q_R}^c & \bar{D}_{Q_R}^{c'} & \bar{D}_{Q_L}^{c'} \\ \bar{Q}_R Q_R & \bar{Q}_R \bar{\Phi} Q_R C & 0 \\ 0 & \bar{Q}_R Q_R & 0 \\ \bar{Q}_L \bar{Q}_R \bar{C} & \bar{Q}_L Q_R \bar{\Phi} & \bar{Q}_L Q_L \end{pmatrix}. \quad (4.10)$$

The mass spectrum becomes $\lambda^{\bar{q}_r+q_r}$, $\lambda^{\bar{q}_r+q_r}$, and $\lambda^{\bar{q}_l+q_l}$.

The mass of the fields Q and $\bar{Q}$ is obtained from the interaction $\bar{Q}_L Q_L$ as $\lambda^{\bar{q}_l+q_l}$. The mass of the fields U^c and $\bar{U}^c$ is obtained from the interaction $\bar{Q}_R Q_R$ as $\lambda^{q_r+\bar{q}_r}$.

By the above argument, the mass spectrum of superheavy particles are determined only by the anomalous $U(1)_A$ charges, so we can examine whether coupling unification is realized or not. Before going to the discussion in the next subsection, we define the reduced mass matrices $\bar{M}_I$ by getting rid of the massless modes from the original mass matrices M_I . The rank of the reduced matrices in our semi-simple model are $\bar{r}_Q = \bar{r}_{U^c} = 1$, $\bar{r}_{E^c} = 2$, $\bar{r}_L = 7$ and $\bar{r}_{D^c} = 3$. It is useful to define the effective anomalous $U(1)_A$ charges:

$$x_I \equiv i + \frac{1}{2}\Delta\phi, \quad \bar{x}_I \equiv \bar{i} - \frac{1}{2}\Delta\phi, \quad (x = l', l^{c'}, d^{c'}, d'), \quad (4.11)$$

$$x_I \equiv i + \Delta c - \frac{1}{2}\Delta\phi, \quad \bar{x}_I \equiv \bar{i} - \Delta c + \frac{1}{2}\Delta\phi, \quad (x = l, d^c), \quad (4.12)$$

$$x_I \equiv i, \quad \bar{x}_I \equiv \bar{i}, \quad (x = q, u^c, e^c), \quad (4.13)$$

where $I = \Phi, C, C', Q_L, Q_R$ ($i = \phi, c, c', q_l, q_r$), $\Delta\phi \equiv \frac{1}{2}(\phi - \bar{\phi})$, and $\Delta c \equiv \frac{1}{2}(c - \bar{c})$. The determinants of the reduced mass matrices are estimated by simple sums of the effective anomalous $U(1)_A$ charges of massive modes:

$$\det \bar{M}_Q = \lambda^{\bar{q}_{Q_L} + q_{Q_L}} = \lambda^{\bar{q}_l + q_l} \quad (4.14)$$

$$\det \bar{M}_{U^c} = \lambda^{\bar{q}_{Q_R} + q_{Q_R}} = \lambda^{\bar{q}_r + q_r} \quad (4.15)$$

$$\det \bar{M}_{E^c} = \lambda^{\bar{e}_{\Phi}^c + e_{\Phi}^c + \bar{e}_{C'}^c + e_{C'}^c} = \lambda^{\bar{\phi} + \phi + \bar{c}' + c'} \quad (4.16)$$

$$\det \bar{M}_{D^c} = \lambda^{d'_{Q_L} + \bar{d}'_{Q_L} + \bar{d}_{Q_R}^c + d_{Q_R}^c + \bar{d}_{Q_R}^{c'} + d_{Q_R}^{c'}} = \lambda^{\bar{q}_l + q_l + 2(\bar{q}_r + q_r)} \quad (4.17)$$

$$\begin{aligned} \det \bar{M}_L &= \lambda^{l'_C + \bar{l}'_{\Phi} + \bar{l}'_{\bar{C}} + l'_{C'} + \bar{l}'_{C'} + \bar{l}_{\bar{C}} + \bar{l}_{C'} + l'_C + \bar{l}'_{\Phi} + \bar{l}'_{\bar{C}} + l'_{C'} + \bar{l}'_{C'} + l_C + l_{C'}} \\ &= \lambda^{3(c + \bar{c} + c' + \bar{c}') + 2\bar{\phi} - \Delta\phi}. \end{aligned} \quad (4.18)$$

Then all the elements of mass matrices are estimated by simple sum of the effective charges of superheavy particles if they are not vanishing, and the determinants of mass matrices are also determined by simple sum of the effective charges. We will use this result in calculating the running gauge couplings.

5 Coupling unification

In this section, we apply the general discussion on the gauge coupling unification in Ref.[3] to our scenario. The pattern of the breaking of the gauge symmetry in our model is as follows. At the scale $\Lambda_\Phi \sim \lambda^{-(\phi+\bar{\phi})/2}$, $SU(3)^3$ is broken into $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$. The $SU(2)_R \times U(1)_{B-L}$ is broken into $U(1)_Y$ at the scale $\Lambda_C \sim \lambda^{-(c+\bar{c})/2}$. We base our analysis on one loop renormalization group equations. The conditions of the gauge coupling unification are given by

$$\alpha_3(\Lambda) = \alpha_2(\Lambda) = \frac{5}{3}\alpha_Y(\Lambda) \equiv \alpha_1(\Lambda), \quad (5.1)$$

where $\alpha_1^{-1}(\Lambda_\Phi > \mu > \Lambda_C) \equiv \frac{3}{5}\alpha_R^{-1}(\Lambda_\Phi > \mu > \Lambda_C) + \frac{2}{5}\alpha_{B-L}^{-1}(\Lambda_\Phi > \mu > \Lambda_C)$, $\alpha_1^{-1}(\mu > \Lambda_\Phi) \equiv \frac{4}{5}\alpha_{3R}^{-1}(\mu > \Lambda_\Phi) + \frac{1}{5}\alpha_{3L}^{-1}(\mu > \Lambda_\Phi)$, and $\alpha_2^{-1}(\mu > \Lambda_\Phi) \equiv \alpha_{3L}^{-1}(\mu > \Lambda_\Phi) + \frac{2}{5}\alpha_{B-L}^{-1}(\mu > \Lambda_C)$. Here $\alpha_X = \frac{g_X^2}{4\pi}$ and the parameters $g_X (X = 3, 3L, 3R, 2, R, B-L, Y)$ are the gauge couplings of $SU(3)_C$, $SU(3)_L$, $SU(3)_R$, $SU(2)_L$, $SU(2)_R$, $U(1)_{B-L}$ and $U(1)_Y$, respectively.

Using the fact that the three gauge couplings of the minimal SUSY standard model (MSSM) meet at the scale $\Lambda_G \sim 2 \times 10^{16}$ GeV, the above conditions for gauge coupling unification are rewritten

$$b_1 \ln \left(\frac{\Lambda}{\Lambda_G} \right) + \Sigma_I \Delta b_{1I} \ln \left(\frac{\Lambda^{\bar{r}_I}}{\det \bar{M}_I} \right) - \frac{12}{5} \ln \left(\frac{\Lambda}{\Lambda_C} \right) - 2 \ln \left(\frac{\Lambda}{\Lambda_\Phi} \right) \quad (5.2)$$

$$= b_2 \ln \left(\frac{\Lambda}{\Lambda_G} \right) + \Sigma_I \Delta b_{2I} \ln \left(\frac{\Lambda^{\bar{r}_I}}{\det \bar{M}_I} \right) - 2 \ln \left(\frac{\Lambda}{\Lambda_\Phi} \right) \quad (5.3)$$

$$= b_3 \ln \left(\frac{\Lambda_A}{\Lambda_G} \right) + \Sigma_I \Delta b_{3I} \ln \left(\frac{\Lambda_A^{\bar{r}_I}}{\det \bar{M}_I} \right), \quad (5.4)$$

where $(b_1, b_2, b_3) = (33/5, 1, -3)$ are the renormalization group coefficients for MSSM and $\Delta b_{aI} (a = 1, 2, 3)$ are the corrections to the coefficients from the massive fields $I = Q + \bar{Q}, U^c + \bar{U}^c, E^c + \bar{E}^c, D^c + \bar{D}^c$, and $L + \bar{L}$. The second last term in Eq. (5.2) is from the breaking $SU(2)_R \times U(1)_{B-L} \rightarrow U(1)_Y$ by the VEV $\langle C \rangle$, and the last terms in Eqs. (5.2), (5.3) are from the breaking $SU(3)_L \times SU(3)_R \rightarrow SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ by the VEV $\langle \Phi \rangle$. Since all the mass matrices and the symmetry breaking scales appearing in the above conditions are determined by the anomalous $U(1)_A$ charges, these conditions can be translated to the constraint on the effective charge and the cutoff scale,

$$\alpha_1(\Lambda) = \alpha_2(\Lambda) \rightarrow \Lambda \sim \Lambda_G \lambda^{-\frac{1}{28}(5\bar{\phi}-\phi+6(c+\bar{c}))}, \quad (5.5)$$

$$\alpha_2(\Lambda) = \alpha_3(\Lambda) \rightarrow \Lambda \sim \Lambda_G \lambda^{\frac{1}{8}(7\bar{\phi}+\phi+6(c+\bar{c}+c'+\bar{c}'))}, \quad (5.6)$$

$$\alpha_1(\Lambda) = \alpha_3(\Lambda) \rightarrow \Lambda \sim \Lambda_G \lambda^{\frac{1}{96}(25\bar{\phi}+19\phi+30(c+\bar{c})+18(c'+\bar{c}'))}. \quad (5.7)$$

A naive calculation leads to the relation between the charges,

$$59\bar{\phi} + 41\phi + 42(c + \bar{c}) + 54(c' + \bar{c}') \sim 0, \quad (5.8)$$

which is difficult to satisfy in our scenario. However, careful calculation shows that gauge coupling unification is possible, though somewhat larger ambiguities of order one coefficients are required than in a simple group unification. Actually with the typical charge assignment in Table I, the coupling unification is realized as in Fig. 1, using the ambiguities of order one coefficients $\lambda \leq y \leq \lambda^{-1}$.




Figure 1: Here we adopt $\lambda = 0.22$, $\alpha_1^{-1}(M_Z) = 59.47$, $\alpha_2^{-1}(M_Z) = 29.81$, $\alpha_3^{-1}(M_Z) = 8.40$, the SUSY breaking scale $m_{SB} \sim 1$ TeV and the anomalous $U(1)_A$ charges: $(\psi_1, \psi_2, \psi_3) = (9/2, 7/2, 3/2)$, $\phi = -3$, $\bar{\phi} = 2$, $c = -3$, $\bar{c} = 1$, $z = -1$, $c' = 0$ and $\bar{c}' = 4$. Using the ambiguities of coefficients $\lambda \leq y \leq \lambda^{-1}$, three gauge couplings meet at around 10^{16} GeV.

Note that the unified gauge coupling can remain in the perturbative region in $SU(3)^3$ model, which is different from the E_6 model. This is because the Higgs sector in $SU(3)^3$ unification is much simpler than that in E_6 unification, since we do not have to introduce adjoint Higgses to realize the doublet-triplet splitting.

The cutoff scale tends to be lower than the Planck scale. Indeed, the cutoff is taken as 10^{16} GeV in Fig. 1. Since the cutoff scale is so low, we have to take care of proton decay via dimension five operators [30], which are obtained from

$$\lambda^{\psi_i + \psi_j + \psi_k + \psi_l + \bar{\phi}} \Psi_i \Psi_j \Psi_k \Psi_l \bar{\Phi} \quad (5.9)$$

by developing the VEV $\langle \bar{\Phi} \rangle \sim \lambda^{-\frac{1}{2}(\phi + \bar{\phi})}$. The coefficients are suppressed not only by the usual small Yukawa factor but also by the suppression factor $\lambda^{4n + \frac{1}{2}(\bar{\phi} - \phi)} =$

$\lambda^{8.5}$. Even if we take the cutoff $\Lambda \sim 10^{16}$ GeV, the ‘effective’ colored Higgs mass is around $\lambda^{-8.5}\Lambda \sim 10^{22}$ GeV, which is much larger than the experimental bound of 10^{18} GeV. Thus, proton decay via dimension five operator is adequately suppressed.

On the other hand, proton decay $p \rightarrow e^+\pi^0$ via dimension six operators from the Kähler potential

$$K = \frac{1}{\Lambda^2} \Psi_1^\dagger \Psi_1 \Psi_1^\dagger \Psi_1, \quad (5.10)$$

which are allowed by the symmetry in our scenario by taking the unification scale Λ_U as the cutoff Λ , may be seen in future experiments. If we roughly estimate the lifetime of proton using the formula in Ref. [32] and the recent result of the lattice calculation for the hadron matrix element parameter α [33], we find

$$\tau_p(p \rightarrow e^+\pi^0) \sim 4.5 \times 10^{34} \left(\frac{\Lambda}{10^{16} \text{GeV}} \right)^4 \left(\frac{0.015(\text{GeV})^3}{\alpha} \right)^2 \text{ years}. \quad (5.11)$$

This estimate, albeit a rough one, provides a strong motivation for continuing the proton decay search.

6 Discussion and Summary

Besides $SU(3)^3$, E_6 has the other maximal semi-simple subgroups $SU(6) \times SU(2)_L$ and $SU(6) \times SU(2)_R$ [35]. The matter sector can be applied to these subgroups in a straightforward way. However, in the Higgs sector, it is difficult to realize the situation in which only one pair of doublet Higgs is massless. It is difficult to make the partner of the doublet Higgs massive, while keeping the latter massless. On the contrary, in $SU(3)^3$ gauge symmetry, since the partners of the doublet Higgs L , E^c and N^c are absorbed by the Higgs mechanism, one pair of doublet Higgs can be massless.

In the typical charge assignment, the charges of the matter sector respect the E_6 symmetry, while those of the Higgs sector do not. It is difficult to respect E_6 symmetry in the Higgs sector without additional massless fields other than the fields in the MSSM.

By introducing singlet fields, we can build models with integer Kac-Moody level. For example, in addition to the fields in Table I, we introduce a singlet field with charge 10, one with charge -8, 43 singlet fields with charge $3/2$, and 62 singlet fields with charge $1/2$. Then using the relation ¹

$$\frac{C_a}{k_a} = \frac{1}{3k_A} \text{tr } Q_A^3 = \frac{1}{24} \text{tr } Q_A, \quad (6.1)$$

¹ $C_a \equiv \text{Tr}_{G_a} T(R) Q_A$. Here $T(R)$ is the Dynkin index of the representation R , and we use the convention in which $T(\text{fundamental rep.}) = 1/2$.

where k_A and k_a are Kac-Moody levels of $U(1)_A$ and $SU(3)_a$ ($a = C, L, R$), these Kac-Moody levels can be calculated as

$$k_A = 4, \quad k_C = k_L = k_R = 2. \quad (6.2)$$

Note that introducing the singlets with charge 10 and -8, the μ problem is solved by the mechanism proposed in Ref. [2].

In our model the difference between the mass matrices of down-type quark and the charged leptons are realized because the matrices are from different Yukawa interactions. However, if this model is regarded as the low energy theory of $E_8 \times E_8$ heterotic superstring theory, we have to break the gauge symmetry E_6 into $SU(3)^3$. Since the matter sector respects E_6 symmetry, it is natural to expect that the Yukawa interactions also respect it. In order to realize the different Yukawa interactions, we have to implement the breaking. In the brane world scenario, there is an interesting mechanism to break the gauge symmetry[9]. However, it seems difficult to realize the breaking in the Yukawa coupling of matter which resides on the brane. To enforce the E_6 breaking some of the matter must be in the bulk, where the E_6 gauge symmetry is not respected.

In this paper, we have proposed a realistic semi-simple unified theory with $SU(3)^3$ gauge group. Since generic interactions have been introduced, we can define the model by the anomalous $U(1)_A$ charges. Large neutrino mixing angles can be realized in the model. Moreover, the FCNC process is automatically suppressed. The half integer charges of the matter sector automatically play the role of R -parity. The model has the same matter structure as in E_6 model[4], but different and simpler Higgs structure. Actually we do not need the adjoint Higgs fields to realize doublet-triplet splitting. And the gauge couplings at the cutoff scale can be in the perturbative region. Note that in the $SU(3)^3$ model, in contrast to $SU(5)$, $SO(10)$ or E_6 , the lightest magnetic monopole carries three (instead of one) quanta of Dirac magnetic charge[36]. This is readily seen by noting that one is allowed, in principle, to include non-bifundamental vector-like representations such as $(\mathbf{1}, \mathbf{3}, \mathbf{1}) + (\mathbf{1}, \bar{\mathbf{3}}, \mathbf{1})$ that, despite their color singlet nature, carry fractional($e/3$) electric charge. The Dirac quantization then requires that the corresponding magnetic charge has three units. The number density of primordial $SU(3)^3$ monopoles depends, of course, on the underlying cosmological scenario, and should not exceed the nominal Parker bound of about $10^{-16} \text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$. The discovery of magnetic monopoles would be a truly remarkable event, and measurement of their magnetic charge would allow us to distinguish between a variety of unified gauge theories.

It would be interesting to extend the approach presented here to other semi-simple unification schemes. For instance, the gauge symmetry $SU(3)^3$ with three $\mathbf{27}$'s of E_6 can be embedded, in principle, in $SU(4) \times SU(3) \times SU(3)$ [37] which could be worth pursuing.

7 Acknowledgement

We would like to thank the organizers of the workshops NOON2001 at Kashiwa and post NOON at Kyoto, where this collaboration began. Q.S. is supported in part by the DOE under Grant No. DE-FG02-91ER40626.

References

- [1] N. Maekawa, Prog. Theor. Phys. **106** (2001) 401 (hep-ph/0104200); hep-ph/0110276.
- [2] N. Maekawa, Phys. Lett. **B521** (2001) 42 (hep-ph/0107313).
- [3] N. Maekawa, Prog. Theor. Phys. **107** (2002) 597 (hep-ph/0111205).
- [4] M. Bando and M. Maekawa, Prog. Theor. Phys. **106** (2001) 1255 (hep-ph/0109018).
- [5] N. Maekawa and T. Yamashita, hep-ph/0202050.
- [6] E. Witten, Phys. Lett. **B149** (1984),351;
M. Dine, N. Seiberg and E. Witten, Nucl. Phys. **B289** (1987), 589;
J.J. Atick, L.J. Dixon and A. Sen, Nucl. Phys. **B292** (1987),109;
M. Dine, I. Ichinose and N. Seiberg, Nucl. Phys. **B293** (1987),253.
- [7] M. Green and J. Schwarz, Phys. Lett. **B149** (1984),117.
- [8] E. Witten, Phys. Lett. **B105** (1981) 267;
A. Masiero, D.V. Nanopoulos, K. Tamvakis and T.Yanagida, Phys. Lett. **115** (1982) 380;
B. Grinstein, Nucl. Phys. **B206** (1982) 387;
K. Inoue, A. Kakuto and T. Takano, Prog. Theor. Phys. **75** (1986) 664;
E. Witten, *Nucl. Phys.* **B258** (1985) 75;
T. Yanagida, Phys. Lett. **B344** (1995) 211;
- [9] Y. Kawamura, Prog. Theor. Phys. **105** (2001) 691;*ibid* **105** (2001) 999;
L. Hall and Y. Nomura, Phys. Rev. **D64** (2001) 055003.
- [10] S. Dimopoulos and F. Wilczek, NSF-ITP-82-07;
M. Srednicki, Nucl. Phys. **B202** (1982) 327.
- [11] S.M. Barr and S. Raby, Phys. Rev. Lett. **79** (1997) 4748.
- [12] Z. Chacko and R.N. Mohapatra, Phys. Rev. **D59** (1999) 011702; Phys. Rev. Lett. **82** (1999) 2836.

- [13] K.S. Babu and S.M. Barr, Phys. Rev. **D48** (1993) 5354;*ibid* **D50** (1994) 3529.
- [14] Fukuda et al.(The Super-Kamiokande Collaboration), Phys. Lett. **B436** (1998) 33; Phys. Rev. Lett. **81** (1998) 1562; Phys. Rev. Lett. **86** (2001) 5656.
- [15] C.D. Froggatt and H.B. Nielsen, Nucl. Phys. **B147** (1979) 277.
- [16] See, for instance, M. Green, J. Schwartz and E. Witten, Superstring Theory (Cambridge University Press, 1987) and references therein.
- [17] G. Lazarides, C. Panagiotakopoulos and Q. Shafi, Phys. Lett. **B315** (1993) 325.
- [18] G. Dvali and Q. Shafi, Phys. Lett. **B326** (1994) 258; **B339** (1994) 241.
- [19] G. Dvali and Q. Shafi, Phys. Lett. **B403** (1997) 65.
- [20] T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and Baryon Number in the Universe*, ed. O. Sawada and A. Sugamoto (KEK report 79-18,1979).
M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, ed. P. van Nieuwenhuizen and D. Z. Freedman (North Holland, Amsterdam, 1979).
- [21] The CHOOZ Collaboration, Phys. Lett. **B420** (1998) 397.
- [22] L. Wolfenstein, Phys. Rev. **D17** (1978) 2369.
S. P. Mikheev and A. Smirnov, Yad. Fiz. **42** (1985), 1441; Nouvo Cim. **9C** (1986) 17.
- [23] S. Dimopoulos and G. F. Giudice, Phys. Lett. **B357** (1995) 573.
- [24] A. G. Cohen, D. B. Kaplan and A. E. Nelson, Phys. Lett. **B388** (1996) 588.
- [25] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, Nucl. Phys. **B477** (1996) 321.
- [26] K. Kurosawa and N. Maekawa, Prog. Theor. Phys. **102** (1999) 121.
- [27] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57** (1986) 961.
R. Barbieri and L. Hall, Phys. Lett. **B338** (1994) 212.
J. Hisano, T. Moroi, K. Tobe, M. Yamaguchi and T. Yanagida, Phys. Lett. **B357** (1995) 579.
J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. **D53** (1996) 2442.
J. Sato and K. Tobe, Phys. Rev. **D63** (2001) 116010.

- [28] M. Drees, Phys. Lett. **B181** (1986) 279.
Y. Kawamura, H. Murayama and M. Yamaguchi, Phys. Lett. **B324** (1994) 52; Phys. Rev. **D51** (1995) 1337.
- [29] M. Bando and T. Kugo, Prog. Theor. Phys. **101** (1999) 1313;
M. Bando, T. Kugo and K. Yoshioka, Prog. Theor. Phys. **104** (2000) 211.
- [30] N. Sakai and T. Yanagida, Nucl. Phys. **B197** (1982) 533.
- [31] T. Goto and T. Nihei, Phys. Rev. **D59** (1999) 115009.
- [32] J. Hisano, H. Murayama and T. Yanagida, Nucl. Phys. **B402** (1993) 46.
- [33] JLQCD Collaboration, S. Aoki et al., Phys. Rev. **D62** (2000) 014506.
- [34] Super-Kamiokande Collaboration, Phys. Rev. Lett. **81** (1998) 3319; *ibid* **83** (1999) 1529.
- [35] M. Matsuda and T. Matsuoka, Phys. Lett. **B487** (2000) 104; *ibid* **B499** (2001) 287.
- [36] Q. Shafi, Proc. of NATO ASI "Monopole '83", Vol. 111, page 47.
- [37] T.W. Kephart and Q. Shafi, Phys. Lett. **B 520** (2001)313 (hep-ph/0105237).